



What is claimed:

1. A compound represented by the Formula I:



wherein:

- 5 R^1 is a moiety represented by the formula



where


Z is selected from the group consisting of CH and NH, and Q is a moiety such that
 10 R^1 is a substituted or unsubstituted monocyclic or bicyclic heteroaryl which has at least two carbon atoms in the heteroaryl ring system;

X is selected from the group consisting of CH_2 , O, S, and NH;

Y is selected from the group consisting of CH_2 , O, and S, provided that at least one of X and Y is CH_2 , or X and Y together with the bond there-between form a cyclopropyl;

15 R^2 and R^3 are independently selected from the group consisting of hydrogen, methyl, halogen, trifluoromethyl, and cyano; and

R^4 is selected from the group consisting of




where R^5 is selected from the group consisting of substituted and unsubstituted
 20 aryl, heteroaryl, cycloalkyl, heterocycloalkyl, $O-R^7$, NR^8R^9 , C_1-C_8 alkyl, and monocyclic heterocycloalkyl, R^6 is selected from the group consisting of substituted and unsubstituted aryl, heteroaryl, cycloalkyl, heterocycloalkyl, alkenyl, $O-R^7$, $C(O)R^7$, NR^8R^9 , C_2-C_8 alkyl, and monocyclic heterocycloalkyl, where R^7 is selected from the group consisting of substituted and unsubstituted alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, R^8
 25 is selected from the group consisting of hydrogen, and substituted and unsubstituted alkyl,

and R⁹ is selected from the group consisting of substituted and unsubstituted alkyl, aryl, heteroaryl, cycloalkyl, and heterocycloalkyl;

or a pharmaceutically acceptable prodrug, pharmaceutically active metabolite, or pharmaceutically acceptable salt thereof.

- 5 2. A compound according to claim 1, wherein R¹ is a substituted or unsubstituted heteroaryl group selected from the group consisting of:




X is selected from the group consisting of CH₂, O, and S;

Y is selected from the group consisting of CH₂ and S, provided that at least one of

- 10 X and Y is CH₂;

R² and R³ are independently selected from the group consisting of hydrogen, methyl, fluorine, and chlorine, and

R⁴ is selected from the group consisting of




- 15 where R⁵ is selected from the group consisting of substituted and unsubstituted aryl, heteroaryl, cycloalkyl, heterocycloalkyl, O-R⁷, NR⁸R⁹, C₁-C₈ alkyl, and monocyclic

heterocycloalkyl, R^6 is selected from the group consisting of substituted and unsubstituted aryl, heteroaryl, cycloalkyl, heterocycloalkyl, alkenyl, $O-R^7$, $C(O)R^7$, NR^8R^9 , C_2-C_8 alkyl, and monocyclic heterocycloalkyl, where R^7 is selected from the group consisting of substituted and unsubstituted alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, R^8 is selected from the group consisting of hydrogen and substituted and unsubstituted alkyl, and R^9 is selected from the group consisting of substituted and unsubstituted alkyl, aryl, heteroaryl, cycloalkyl, and heterocycloalkyl;

or a pharmaceutically acceptable prodrug, pharmaceutically active metabolite, or pharmaceutically acceptable salt thereof.

3. A compound represented by the Formula II:




wherein:

X is selected from the group consisting of CH_2 , O, and S;

Y is selected from the group consisting of CH_2 and S, provided that at least one of X and Y is CH_2 ;

R^2 and R^3 are independently selected from the group consisting of hydrogen, methyl, fluorine, and chlorine;


R^4 is selected from the group consisting of



where R^5 and R^6 are each independently selected from the group consisting of substituted and unsubstituted aryl and heteroaryl; and R^{10} is selected from the group consisting of substituted and unsubstituted alkenyl, aryl, heteroaryl, and HNR^9 , where R^9 is selected from the group consisting of substituted and unsubstituted alkyl, aryl, heteroaryl, cycloalkyl, and heterocycloalkyl;

or a pharmaceutically acceptable prodrug, pharmaceutically active metabolite, or pharmaceutically acceptable salt thereof.

4. A compound according to claim 3, wherein R^5 and R^6 are each independently selected from the group consisting of substituted and unsubstituted aryl; or a pharmaceutically acceptable salt, a pharmaceutically acceptable prodrug, or a pharmaceutically active metabolite thereof, or a pharmaceutically acceptable salt of said metabolite.
5. A compound according to claim 3, wherein R^5 and R^6 are each independently selected from the group consisting of substituted and unsubstituted heteroaryl; or a pharmaceutically acceptable salt, a pharmaceutically acceptable prodrug, or a pharmaceutically active metabolite thereof, or a pharmaceutically acceptable salt of said metabolite.
6. A compound represented by the Formula III:




wherein:

X is selected from the group consisting of CH_2 , O, S, and NH;

Y is selected from the group consisting of CH_2 , O, and S, provided that at least one of X and Y is CH_2 , or X and Y together with the bond there-between form a cyclopropyl;

R^2 and R^3 are independently selected from the group consisting of hydrogen, methyl, halogen, trifluoromethyl, and cyano; and

R^4 is selected from the group consisting of



where R^5 is selected from the group consisting of substituted and unsubstituted aryl, heteroaryl, cycloalkyl, heterocycloalkyl, $O-R^7$, NR^8R^9 , C_1-C_8 alkyl, and monocyclic heterocycloalkyl, R^6 is selected from the group consisting of substituted and unsubstituted

aryl, heteroaryl, cycloalkyl, heterocycloalkyl, alkenyl, $O-R^7$, $C(O)R^7$, NR^8R^9 , C_2-C_8 alkyl, and monocyclic heterocycloalkyl, where R^7 is selected from the group consisting of substituted and unsubstituted alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl, R^8 is selected from the group consisting of hydrogen and substituted and unsubstituted alkyl, and R^9 is selected from the group consisting of substituted and unsubstituted alkyl, aryl, heteroaryl, cycloalkyl, and heterocycloalkyl;

or a pharmaceutically acceptable prodrug, pharmaceutically active metabolite, or pharmaceutically acceptable salt thereof.


7. A compound according to claim 6, wherein:

X is selected from the group consisting of CH_2 , O, and S;

Y is selected from the group consisting of CH_2 and S, provided that at least one of X and Y is CH_2 ;

R^2 and R^3 are independently selected from the group consisting of hydrogen, methyl, fluorine, and chlorine; and

R^4 is selected from the group consisting of



where R^5 and R^6 are each independently selected from the group consisting of substituted and unsubstituted aryl and heteroaryl;

or a pharmaceutically acceptable salt, a pharmaceutically acceptable prodrug, or a pharmaceutically active metabolite thereof, or a pharmaceutically acceptable salt of said metabolite.

8. A compound according to claim 7, wherein R^5 and R^6 are each independently selected from the group consisting of substituted and unsubstituted aryl; or a pharmaceutically acceptable salt, a pharmaceutically acceptable prodrug, or a pharmaceutically active metabolite thereof, or a pharmaceutically acceptable salt of said metabolite.

9. A compound according to claim 7, wherein R^5 and R^6 are each independently selected from the group consisting of substituted and unsubstituted heteroaryl; or a pharmaceutically acceptable salt, a pharmaceutically acceptable prodrug,

or a pharmaceutically active metabolite thereof, or a pharmaceutically acceptable salt of said metabolite.


10. A compound according to claim 7, wherein:

X is CH₂;

Y is S;

R² and R³ are independently selected from the group consisting of hydrogen, methyl, fluorine, and chlorine; and

R⁴ is selected from the group consisting of



where R⁵ and R⁶ are each independently selected from the group consisting of substituted and unsubstituted aryl and heteroaryl;

or a pharmaceutically acceptable salt, a pharmaceutically acceptable prodrug, or a pharmaceutically active metabolite thereof, or a pharmaceutically acceptable salt of said metabolite.

11. A compound according to claim 10, wherein R⁵ and R⁶ are each independently selected from the group consisting of substituted and unsubstituted aryl; or a pharmaceutically acceptable salt, a pharmaceutically acceptable prodrug, or a pharmaceutically active metabolite thereof, or a pharmaceutically acceptable salt of said metabolite.

12. A compound selected from the group consisting of

N-(3,4,5-Trimethoxyphenyl)-3-[(pyrazin-2-yl)sulfanylmethyl]benzamide;

N-(3,4,5-Trimethoxyphenyl)-3-[(5-amino-2H-[1,2,4]triazol-3-yl)sulfanylmethyl]benzamide;

N-(4-Isopropyl-3-methylphenyl)-3-[(pyrazin-2-yl)sulfanylmethyl]benzamide;

N-(4-Isopropyl-3-methylphenyl)-3-[(5-amino-2H-[1,2,4]triazol-3-yl)sulfanylmethyl]benzamide;

N-(4-Isopropyl-3-methylphenyl)-3-[(1H-pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]benzamide;

N-(2-Methylquinolin-6-yl)-3-[(pyrazin-2-yl)sulfanylmethyl]benzamide;

N-(3-Isopropylphenyl)-3-[(pyrazin-2-yl)sulfanylmethyl]benzamide;

N-(3,5-Dibromo-4-methylphenyl)-3-[(pyrazin-2-yl)sulfanylmethyl]benzamide;

N-(3,4,5-Trimethoxyphenyl)-3-[(1H-pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl] benzamide;

N-(3,4,5-Trimethoxyphenyl)-3-[(1H-pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl] benzamide;

N-(Quinolin-6-yl)-3-[(1H-pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]benzamide;

N-(5-Methylisoxazol-3-yl)-3-[(1H-pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl] benzamide;

N-(Pyridin-4-yl)methyl-3-[(1H-pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl] benzamide;

N-(1,3-Benzodioxyl-5-ylmethyl)-3-[(1H-pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl] benzamide;

N-(2-Methoxybenzyl)-3-[(1H-pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]benzamide;

N-(2-Phenylethyl)-3-[(1H-pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]benzamide;

N-(2-Methoxyphenyl)-3-[(1H-pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]benzamide;

N-[3-(N-Methyl-N-phenylamino)propyl]-3-[(5-methyl-1H-1,2,4-triazol-3-yl)sulfanylmethyl]benzamide;

N-(1,3-Benzodioxyl-5-ylmethyl)-3-[(5-methyl-1H-1,2,4-triazol-3-yl)sulfanylmethyl] benzamide;

N-[4-cyano-3-(trifluoromethyl)phenyl]-3-[(1H-pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]benzamide;

N-(3,3-Diphenylpropyl)-3-[(5-methyl-1H-1,2,4-triazol-3-yl)sulfanylmethyl]benzamide;

3-[(5-Methyl-1H-1,2,4-triazol-3-yl)sulfonylmethyl]-N-phenethylbenzamide;

3-[(1H-Pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]-N(3-isopropylphenyl)benzamide;

3-[(1H-Pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]-N(3-trifluoromethyl-5-methoxyphenyl)-benzamide;

3-[(1H-Pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]-N(3,5-bis-trifluoromethylphenyl)-benzamide;

3-[(1H-Pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]-N(3-t-butylphenyl)-benzamide;

5 3-[(1H-Pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]-N(4-isopropylphenyl)-benzamide;

3-[(1H-Pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]-N(4-trifluoromethoxyphenyl)-benzamide;

10 3-[(1H-Pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]-N(3,5-dimethylphenyl)-benzamide;

3-[(1H-Pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]-N(3-(2-hydroxyethyl)phenyl)-benzamide;

3-[(1H-Pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]-N(4-dimethylaminophenyl)-benzamide;

15 3-[(1H-Pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]-N(3-trifluoromethylsulfonyl phenyl)-benzamide;

3-[(1H-Pyrazolo[3,4-d]pyrimidin-4-yl)sulfanylmethyl]-N(3-dimethylaminophenyl)-benzamide;

20 3-[(5-Cyanoamino-2H-[1,2,4]triazol-3-yl)sulfanylmethyl]-N-(3,4,5-trimethoxyphenyl) benzamide;

3-[(5-(Methoxycarbonylamino)-2H-[1,2,4]triazol-3-yl)sulfanylmethyl]-N-(3,4,5-trimethoxyphenyl)benzamide;

N-(3,4,5-Trimethoxyphenyl)-3-[(5-acetylamino-2H-[1,2,4]triazol-3-yl)sulfanylmethyl] benzamide;

25 N-(4-Isopropyl-3-methylphenyl)-3-[(pyrazin-2-yl)methylsulfanyl]benzamide;

N-(2-Methylquinolin-6-yl)-3-[(pyrazin-2-yl)methylsulfanyl]benzamide;

N-(2-Methyl-quinolin-6-yl)-3-(pyridin-3-ylmethylsulfanyl)-benzamide dihydrochloride;

30 N-(2-methyl-quinolin-6-yl)-3-[[5-(phenylamino)-2-H-pyrazol-3-yl)methylsulfanyl] benzamide;

N-(3,4,5-trimethoxyphenyl)-3-[2-(5-phenylamino-2H-pyrazol-3-yl)ethyl]benzamide;

3-[[5-((E)-2-(4-Hydroxy-3-methoxyphenyl)ethenyl)-2H-pyrazol-3-yl]-methylsulfanyl]-N-(2-methylquinolin-6-yl)benzamide;

3-[5-(2-(3,4-Dimethoxyphenyl)ethenyl)-2H-pyrazol-3-yl)methylsulfanyl]-N-(2-methylquinolin-6-yl)benzamide;

5 3-(2-{5-[(E)-2-(3,4-Dimethoxyphenyl)ethenyl]-2H-pyrazol-3-yl}-ethyl)-N-(3-methyl-4-isopropylphenyl)-benzamide;

4-Fluoro-3-[[5-((E)-1-propenyl)-2H-pyrazol-3-yl]methoxy]-N-[4-(pyrrolidin-1-yl)-3-trifluoromethylphenyl]benzamide;

10 3-(2-{5-[(E)-2-(3,4-Dimethoxyphenyl)ethenyl]-2H-pyrazol-3-yl}-ethyl)-N-(3-methyl-4-isopropylphenyl)-benzamide;

N-(4-Isopropyl-3-methylphenyl)-3-{2-[5-(4-(methylsulfamoyl)-phenylamino)-2H-pyrazol-3-yl]-ethyl}-benzamide;

N-(2-Methylquinolin-6-yl)-3-[2-(5-phenylamino-2H-pyrazol-3-yl)ethyl]benzamide;

15 N-(4-isopropyl-3-methylphenyl)-3-[2-(5-phenylamino-2H-pyrazol-3-yl)ethyl]benzamide;

N-(4-Isopropyl-3-methylphenyl)-3-{2-[5-(6-methoxypyridin-3-yl)amino-2H-pyrazol-3-yl]-ethyl}-benzamide;

20 N-(4-Dimethylamino-3-trifluoromethylphenyl)-3-{2-[5-(6-methoxypyridin-3-yl)amino-2H-pyrazol-3-yl]ethyl}-benzamide;

N-(6-Dimethylamino-5-trifluoromethylpyridin-3-yl)-3-{2-[5-(6-methoxypyridin-3-yl)amino-2H-pyrazol-3-yl]ethyl}-benzamide;

N-(3,5-Dichloro-4-dimethylaminophenyl)-3-{2-[5-(6-methoxy-pyridin-3-yl)amino-2H-pyrazol-3-yl]ethyl}benzamide;

25 3-{2-[5-(6-Methoxypyridin-3-yl)amino-2H-pyrazol-3-yl]-ethyl}-N-(4-pyrrolidin-1-yl-3-trifluoromethylphenyl)benzamide;

3-{2-[5-(6-Methoxypyridin-3-yl)amino-2H-pyrazol-3-yl]-ethyl}-N-[4-(4-*t*-butoxycarbonylpiperazin-1-yl)-3-trifluoromethylphenyl]benzamide;

30 3-{2-[5-(6-Methoxypyridin-3-yl)amino]-2H-pyrazol-3-yl]ethyl}-N-(4-piperazin-1-yl-3-trifluoromethylphenyl)benzamide;

4-Fluoro-3-[[5-(pyridin-3-yl)amino-2H-pyrazol-3-yl]methoxy]-N-[[4-(pyrrolidin-1-yl)-3-trifluoromethylphenyl]benzamide;

N-(4-Isopropyl-3-methyl-phenyl)-3-[2-(5-phenylamino-2-H-pyrazol-3-yl)-cyclopropyl]-benzamide;

3-[(3-[(E)-2-(4-hydroxy-3-methoxyphenyl)ethenyl]-1H-pyrazol-5-yl)methylamino]-N-(3-methyl-4-isopropylphenyl)benzamide;

5 3-[(5-[(E)-2-(4-hydroxy-3-methoxyphenyl)ethenyl]-1H-pyrazol-3-yl)methylamino]-N-phenyl)benzamide;

4-Fluoro-N-[4-(imidazol-1-yl)-3-trifluoromethylphenyl]-3-[5-(6-methoxypyridin-3-yl)amino-2H-pyrazol-3-ylmethoxy]-benzamide;

10 4-Fluoro-3-[5-(6-methoxy-pyridin-3-yl)amino-2H-pyrazol-3-yl]methoxy-N-(4-pyrrolidin-1-yl-3-trifluoromethyl-phenyl)-benzamide;

4-Fluoro-3-[5-(6-methoxypyridin-3-yl)amino-2H-pyrazol-3-yl]methoxy-N-(3-methoxy-5-trifluoromethyl-phenyl)-benzamide;

N-(4-Isopropyl-3-methyl-phenyl)-3-(Isoquinolin-4-yl)methoxy-benzamide;

15 3-(Isoquinolin-4-yl)methoxy-N-(3,4,5-trimethoxyphenyl)benzamide hydrochloride;

3-(Isoquinolin-4-yl)methoxy-N-(2-methyl-quinolin-6-yl)-benzamide hydrochloride;

3-(Isoquinolin-4-yl)methoxy-N-(2-methyl-4-methylsulfanyl-quinolin-6-yl)-benzamide hydrochloride;

20 3-(Pyridin-3-yl)methoxy-N-(3,4,5-trimethoxyphenyl)benzamide;

N-(Naphthalen-2-yl)-3-(pyridin-3-yl)methoxybenzamide;

N-(1-Allyl-1H-indol-5-yl)-3-(pyridin-3-yl)methoxy-benzamide;

3-(Pyridin-3-yl)methoxy-N-quinolin-6-yl-benzamide;

N-(2-Methyl-quinolin-6-yl)-3-(pyridin-3-yl)methoxy-benzamide;

25 N-(4-Isopropyl-3-methyl-phenyl)-4-fluoro-3-(Isoquinolin-4-yl)methoxy-benzamide;

N-(4-Isopropyl-3-methyl-phenyl)-4-methyl-3-(Isoquinolin-4-yl)methoxy-benzamide;

N-(4-Isopropyl-3-methyl-phenyl)-4-chloro-3-(Isoquinolin-4-yl)methoxy-benzamide;

3-(6-Aminopyridin-3-yl)methoxy-N-(4-Isopropyl-3-methyl-phenyl)benzamide;

3-(6-Aminopyridin-3-yl)methoxy-N-(2-methyl-quinolin-6-yl)-benzamide;

30 3-(6-Acetylaminopyridin-3-yl)methoxy-N-(2-methyl-quinolin-6-yl)-benzamide;

3-(6-Acetylaminopyridin-3-yl)methoxy-N-(4-isopropyl-3-methyl-phenyl)-benzamide;

4-Fluoro-N-(1,2,3,4-tetrahydroquinolin-6-yl)-3-(isoquinolin-4-yl-methoxy)-
benzamide bistrifluoroacetic acid salt;

N-(2,2-difluorobenzo[1,3]dioxol-4-yl-ethyl)-benzamide trifluoroacetic acid salt;

4-Fluoro-N-(2-methyl-1,2,3,4-tetrahydroquinolin-6-yl)-3-(isoquinolin-4-yl-methoxy)-
5 benzamide bistrifluoroacetic acid salt;

N'-{4-[3-(4-Isopropyl-3-methyl-phenylcarbamoyl)-phenoxy];

N-(4-Isopropyl-3-methyl-phenyl)-3-{1-[N'-(3-methoxy-benzylidene)-hydrazino]-
isoquinolin-4-ylmethoxy}-benzamide;

N-(3,5-Diallyl-4-methyl-phenyl)-3-(isoquinolin-4-ylmethoxy)-benzamide;

10 N-(3,5-Dibromo-4-methyl-phenyl)-3-(isoquinolin-4-ylmethoxy)-benzamide;

3-(Isoquinolin-4-ylmethoxy)-N-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-naphthalen-2-
yl)-benzamide;

3-(Isoquinolin-4-ylmethoxy)-N-(3-trifluoromethoxy-phenyl)-benzamide;

N-(2,4-Dimethylquinolin-6-yl)-3-(isoquinolin-4-ylmethoxy)-benzamide;

15 3-(Isoquinolin-4-ylmethoxy)-benzoic acid N'-(4-trifluoromethyl-phenyl)-hydrazide;

N-Benzylloxy-3-(isoquinolin-4-ylmethoxy)-benzamide;

3-(Isoquinolin-4-ylmethoxy)-benzoic acid N'-phenyl-hydrazide;

N-(5,7-dimethyl[1,8]naphthydrin-2-yl)-3-(isoquinolin-4-ylmethoxy)-benzamide;

3-(Isoquinolin-4-ylmethoxy)-N-(1,1,3,3-tetramethyl-1,3-dihydroisobenzofuran-5-yl)-

20 benzamide;

N-(3,5-Dichloro-4-pyrrolidin-1-yl-phenyl)-4-fluoro-3-(pyridin-3-ylmethoxy)-
benzamide;

4-Fluoro-N-(4-morpholin-4-yl-3-trifluoromethyl-phenyl)-3-(pyridin-3-ylmethoxy)-
benzamide;

25 4-Fluoro-N-[4-(piperazin-1-yl)-3-trifluoromethylphenyl]-3-(pyridin-3-
yl)methoxybenzamide;

4-Fluoro-N-(4-morpholin-4-yl-3-trifluoromethyl-phenyl)-3-(isoquinolin-4-
ylmethoxy)-benzamide;

4-Fluoro-N-(4-piperazin-1-yl-3-trifluoromethyl-phenyl)-3-(isoquinolin-4-ylmethoxy)-
30 benzamide;

4-Fluoro-N-(4-morpholin-4-yl-3-trifluoromethyl-phenyl)-3-(quinolin-3-ylmethoxy)-
benzamide;

4-Fluoro-N-(4-piperazin-1-yl-3-trifluoromethyl-phenyl)-3-(quinolin-3-ylmethoxy)-benzamide;

N-(3,5-Dichloro-4-morpholin-4-yl-phenyl)-4-fluoro-3-(pyridin-3-ylmethoxy)-benzamide;

5 N-(3,5-Dichloro-4-piperazin-1-yl-phenyl)-4-fluoro-3-(pyridin-3-ylmethoxy)-benzamide;

4-Fluoro-N-[4-(piperazin-1-yl)-3-trifluoromethylphenyl]-3-(pyridin-3-yl)methoxybenzamide;

10 4-Fluoro-N-(4-(imidazol-1-yl-3-trifluoromethylphenyl)-3-(pyridin-3-yl)methoxybenzamide;

4-Fluoro-N-(4-pyrazol-1-yl-3-trifluoromethyl-phenyl)-3-(pyridin-3-ylmethoxy)-benzamide;

4-Fluoro-3-(pyridin-3-ylmethoxy)-N-(4-[1,2,4]triazol-1-yl-3-trifluoromethyl-phenyl)-benzamide;

15 N-(3,5-Dichloro-4-imidazol-1-yl-phenyl)-4-fluoro-3-(pyridin-3-ylmethoxy)-benzamide;

3-(5-Bromo-pyridin-3-ylmethoxy)-4-fluoro-N-(4-piperazin-1-yl-3-trifluoromethyl-phenyl)-benzamide;

3-(2-Isoquinolin-4-yl-ethyl)-N-phenyl-benzamide;

20 3-(2-Isoquinolin-4-yl-ethyl)-N-(3,3,5-trimethyl-cyclohexyl)-benzamide;

N-(4-Isopropyl-3-methyl-phenyl)-3-(2-isoquinolin-4-yl-ethyl)-benzamide;

3-(2-Isoquinolin-4-yl-ethyl)-N-(2-methyl-quinolin-6-yl)-benzamide;

N-(3,5-Dibromo-4-methyl-phenyl)-3-(2-isoquinolin-4-yl-ethyl)-benzamide;

N-(4,6-Dimethyl-pyridin-2-yl)-3-(2-isoquinolin-4-yl-ethyl)-benzamide;

25 2-Chloro-4-fluoro-N-(4-isopropyl-3-methyl-phenyl)-5-(2-isoquinolin-4-yl-ethyl)-benzamide;

2,4-Difluoro-N-(4-isopropyl-3-methyl-phenyl)-5-(2-isoquinolin-4-yl-ethyl)-benzamide;

2-Fluoro-N-(4-isopropyl-3-methyl-phenyl)-5-(2-isoquinolin-4-yl-ethyl)-benzamide;

30 N-(2-Methyl-quinolin-6-yl)-3-(2-pyridin-3-yl-ethyl)-benzamide hydrochloride;

N-(4-Isopropyl-3-methyl-phenyl)-3-(2-pyridin-3-yl-ethyl)-benzamide;

- N-{3-[(1H-pyrazolo[3,4-d]-pyrimidin-4-ylsulfanylmethyl]phenyl}-(3-bromo-4-methyl) benzamide;
- N-{3-[(1H-pyrazolo[3,4-d]-pyrimidin-4-ylsulfanylmethyl]phenyl}-3,5-bis(trifluoromethyl) benzamide;
- 5 N-{3-[(1H-pyrazolo[3,4-d]-pyrimidin-4-yl)sulfanylmethyl]phenyl}-(4-hydroxy-3-methoxy) benzamide;
- N-{3-[(1H-pyrazolo[3,4-d]-pyrimidin-4-yl)sulfanylmethyl]phenyl}-(4-hydroxy-3-t-butyl) benzamide;
- N-{3-[(1H-pyrazolo[3,4-d]-pyrimidin-4-yl)sulfanylmethyl]phenyl}-4-t-
- 10 butylbenzamide;
- N-{3-[(1H-pyrazolo[3,4-d]-pyrimidin-4-yl)sulfanylmethyl]phenyl}-(4-phenoxy)benzamide;
- N-{3-[(1H-pyrazolo[3,4-d]pyrimidin-4-ylsulfanyl)methyl]phenyl}-N'-[3,5-bis-(trifluoromethyl)phenyl]urea;
- 15 N-{3-[(1H-pyrazolo[3,4-d]pyrimidin-4-ylsulfanyl)methyl]phenyl}-N'-(pyridin-3-yl)urea;
- N-{3-[(1H-pyrazolo[3,4-d]-pyrimidin-4-yl)sulfanylmethyl]phenyl}-(3,5-di-t-butyl) benzamide;
- 3-Bromo-4-hydroxy-N-{3-[(1H-pyrazolo[3,4-d]-pyrimidin-4-
- 20 yl)sulfanylmethyl]phenyl}-benzamide;
- N-{3-[(1H-pyrazolo[3,4-d]-pyrimidin-4-yl)sulfanylmethyl]phenyl}-quinoline-6-carboxamide;
- 5-Fluoro-N-{3-[(1H-pyrazolo[3,4-d]-pyrimidin-4-yl)sulfanylmethyl]phenyl}-indole-2-carboxamide;
- 25 N-{3-[(1H-pyrazolo[3,4-d]-pyrimidin-4-yl)sulfanylmethyl]phenyl}-indole-6-carboxamide;
- (R/S)-2-(2-methylphenyl)-N-{3-[(1H-pyrazolo[3,4-d]pyrimidin-4-ylsulfanyl)methyl]-methyl} phenyl} butanamide;
- 3-*t*-Butyl-4-hydroxy-N-{3-[5-(6-methoxy-pyridin-3-ylamino)-2 H-pyrazol-3-ylmethyl
- 30 sulfanyl]-phenyl}-benzamide;
- 3-*t*-Butyl-4-hydroxy-N-[3-(pyridin-3-ylmethylsulfanyl)-phenyl]-benzamide;
- 3-*t*-Butyl-4-hydroxy-N-[3-(isoquinolin-4-ylmethylsulfanyl)-phenyl]-benzamide;

N-[3-(5-Bromo-pyridin-3-ylmethoxy)-phenyl]-3-t-butyl-4-hydroxy-benzamide;
 4-Acetoxy-3-t-butyl-N-[3-(pyridin-3-ylmethoxy)phenyl]-benzamide;
 4-Acetoxy-3-t-butyl-N-[3-(isoquinolin-4-ylmethoxy)phenyl]-benzamide;
 3-t-Butyl-4-hydroxy-N-[3-(pyridin-3-ylmethoxy)-phenyl]-benzamide;
 5 3-t-Butyl-4-hydroxy-N-[3-(isoquinolin-4-ylmethoxy)-phenyl]-benzamide;
 1-[3-(pyridin-3-ylmethoxy)phenylcarbamoyl]pyrrolidine;
 4-[3-(pyridin-3-ylmethoxy)phenylcarbamoyl]morpholine;
 3-[[6-Methoxy-7-(2-methoxyethoxy)cinnolin-4-yl]sulfanylmethyl]-N-phenyl-
 benzamide;
 10 3-[2-(6-Acetylamino-pyridin-3-yl)-ethyl]-N-(4-piperazin-1-yl-3-trifluoro-
 methylphenyl)-benzamide dihydrochloride;
 3-[2-(6-Amino-pyridin-3-yl)-ethyl]-N-(4-piperazin-1-yl-3-trifluoromethyl-phenyl)-
 benzamide dihydrochloride;
 3-[2-(3H-Imidazo[4,5-b]pyridin-6-yl)-ethyl]-N-(4-piperazin-1-yl-3-trifluoromethyl-
 15 phenyl)-benzamide dihydrochloride;
 5-{2-[3-(Piperazin-1-yl-trifluoromethyl-phenylcarbamoyl)-phenyl]-ethyl}-
 nicotinamide dihydrochloride;
 5-{2-[3-(Piperazin-1-yl-trifluoromethyl-phenylcarbamoyl)-phenyl]-ethyl}-nicotinic
 acid methyl ester dihydrochloride;
 20 4-Fluoro-3-[2-(3H-imidazo[4,5-b]pyridin-6-yl)-ethyl]-N-(4-piperazin-1-yl-3-
 trifluoromethyl-phenyl)-benzamide dihydrochloride; and
 4-Fluoro-3-(5-furan-2-yl-pyridin-3-ylmethoxy)-N-(4-piperazin-1-yl-3-
 trifluoromethyl-phenyl)-benzamide dihydrochloride;
 or a pharmaceutically acceptable salt, pharmaceutically acceptable prodrug, or
 25 pharmaceutically active metabolite thereof or a pharmaceutically acceptable salt of said
 metabolite.

13. A compound selected from the group consisting of the compounds
 corresponding to Example B-27 (Compounds 1-244), Example V-6d
 (Compounds 1-176), Example V-7b (Compounds 1-43) and Example V-14
 30 (Compounds 1-88), or a pharmaceutically acceptable salt, pharmaceutically
 acceptable prodrug, or pharmaceutically active metabolite thereof or a
 pharmaceutically acceptable salt of said metabolite.

14. A pharmaceutically acceptable salt of a pharmaceutically active metabolite of a compound according to claim 1.

15. A pharmaceutical composition for modulating or inhibiting the activity of a protein kinase receptor comprising:

(a) a therapeutically effective amount of an agent selected from the group consisting of a compound according to claim 1, a pharmaceutically acceptable prodrug thereof, a pharmaceutically active metabolite thereof, and a pharmaceutically acceptable salt thereof; and

(b) a pharmaceutically acceptable carrier, diluent, or vehicle therefor.

16. A pharmaceutical composition for modulating or inhibiting the activity of a protein kinase receptor comprising:

(a) a therapeutically effective amount of a pharmaceutically acceptable salt of a pharmaceutically active metabolite of a compound according to claim 1;

(b) a pharmaceutically acceptable carrier, diluent, or vehicle therefor.

17. A method of treating a mammalian disease condition mediated by protein kinase activity, comprising administering to a mammal in need thereof a therapeutically effective amount of an agent selected from the group consisting of a compound according to claim 1, a pharmaceutically acceptable prodrug thereof, a pharmaceutically active metabolite thereof, and a pharmaceutically acceptable salt thereof.

18. A method according to claim 17, wherein the mammalian disease condition is associated with tumor growth, cell proliferation, or angiogenesis.

19. A method of modulating or inhibiting the activity of a protein kinase receptor, comprising contacting the kinase receptor with an effective amount of an agent selected from the group consisting of a compound according to claim 1, a pharmaceutically acceptable prodrug thereof, a pharmaceutically active metabolite thereof, and a pharmaceutically acceptable salt thereof.

20. A method according to claim 19, wherein the protein kinase receptor is a VEGF receptor.